

Investigating the Relationship Between Self-Esteem and Body Image Concern and Self-Compassion in Women with Female Pattern Hair Loss

Azam Bayani¹, Masoud Pourhossein², Mohammad Bayani³

1. Bachelor of Midwifery, Gonabad University of Medical Sciences, Iran.
2. Bachelor of Nursing, Islamic Azad University, Ferdous Branch, Iran.
3. Master's degree in clinical psychology, Islamic Azad University, Torbetjam branch, Iran.

ARTICLE INFO

Keywords:

*Self-Esteem,
Body Image
Concern,
Self-
Compassion,
Hair Loss.*

ABSTRACT

The aim of the present study was to investigate the relationship between self-esteem body image concern and self-compassion in women with hair loss with a female pattern, in this study descriptive-correlation method was used, and the statistical population of this study includes all women with hair loss. It was with a female pattern that they referred to the Runic Skin and Hair Clinic in Qazvin City in the spring of 1402, and the statistical sample of the present study included 300 women with female pattern hair loss who were selected by available sampling. Rosenberg self-esteem scale (RSES), body image concern questionnaire (BICI) and self-compassion scale short form (SCS-SF) were used to collect data. After distributing and completing the questionnaires, the samples were collected and scored. The results showed that self-esteem has a negative and statistically significant relationship with concern about body image in women with hair loss with a feminine pattern, and self-esteem has a statistically significant positive relationship with self-compassion in women with hair loss with a feminine pattern (05/05). 0>P).

1. Introduction

Female androgenetic alopecia or female pattern hair loss is one of the most common types of hair loss in women (Müller et al., 2023). FPHL is characterized by the progressive shrinking of hair follicles and loss of hair density, mainly in the central parietal area of the scalp (Ramos et al., 2015). FPHL is a common hair loss disorder, but with age, 55% of women over 70 years of age experience significant pattern hair loss, and severe disease progression occurs during adolescence in a small number of patients. Typically, FPHL reaches a first peak during the reproductive years and a second peak after menopause (Bertoli et al., 2020). In a Brazilian study, it was found that 68% of women between the ages of 20 and 29 have hair loss (Tsutsui et al., 2022; Chaikittisilpa et al., 2022). reported the prevalence of female pattern hair loss of 2.52% in postmenopausal women with an average age of 58 years. Interestingly, the prevalence of FPHL also varies by location and ethnicity. For example, a prevalence analysis of a student population showed that the overall prevalence of FPHL among high school girls was 28.6%, which was significantly higher in rural than urban areas and significantly associated with family history (Youssef et al., 2022).

On the other hand, androgenetic alopecia can have a significant effect on a person's appearance and self-esteem (Kranz et al., 2019). For women, thick and healthy hair signifies a sense of self-esteem, and self-confidence, shows their ability to change, and security, and implies social interaction. These elements, which are associated with the chronicity of the disease and its treatment resistance, have an important negative impact on the quality of life of women (Shimizu et al., 2018). The results of Bayani et al.'s research (1402) showed that the self-esteem of candidates and non-candidates for hair mesotherapy has a statistically significant difference. Having long and thick hair results in a more beautiful effect for women, the issue of hair loss in can bring more destructive psychological consequences to them, among these consequences can be the loss of self-esteem in women suffering from hair loss. hair, because they feel that they have lost their beautiful appearance by losing a large part of their hair (Bayani et al., 1402). Self-esteem arises from the sense of value and confidence that a person has in himself and his abilities, therefore, self-esteem is a type of personal evaluation that causes a person to pay attention to himself (Ravary and Baldwin, 2018).

Hair loss not only affects aesthetics but also strongly affects the psychological state of people (Alessandrini et al., 2021). Extensive scientific research shows that alopecia affects patients' perception of their body image and significantly reduces their self-confidence (Masood, 2023). Patients who have changes in their physical appearance experience many cognitive changes, including dissatisfaction with their appearance, running away from themselves, and negative thoughts towards themselves (Avis et al., 2004). In this regard, Ayat-Elahi et al. (1401) show that hair loss may lead to dissatisfaction with appearance. Also, the results of Adil and Godwin's research (Adil and Godwin, 2017) indicate that androgenetic alopecia creates a feeling of decreased attractiveness in patients. Body image is actually a person's perception of the size and proportion of the body, which is associated with other people's thoughts, feelings, and views of his body. The term body image refers to people's cognitive evaluation of their bodies (Banfield et al., 2002).

Another variable that can be investigated in women suffering from hair loss with the feminine pattern is self-compassion. (Neff, 2003) defined self-compassion as self-kindness shared humanity, and mindfulness. He introduced the concept of facing one's own personal pain, not avoiding or disconnecting from it, and developing a desire to alleviate this suffering and heal oneself with kindness. Self-compassion or self-love helps adult women to accept more of their bodies, and together with other components of self-compassion, they create perceptions of themselves and their physique for the individual in relation to others, which improve the body image relationship. And confidence helps. In this way, women who love themselves value themselves more (Pisitsungkagarn et al., 2014). Self-compassion may increase women's ability to appreciate their bodies, thereby promoting body acceptance and respect regardless of their imperfections (Stapleton et al., 2017). Hair loss has many psychological and social effects for affected people, especially women, since this problem is not malignant, its annoying conditions are ignored. Therefore, examining the psychological aspects of hair loss in women can bring psychological and physical care for women,

based on this, the present study aims to investigate the relationship between self-esteem and self-compassion and concern about body image in women with hair loss. The hair was done with a feminine pattern.

theories

- Self-esteem is related to body image concerns in women with female-pattern hair loss.
- Self-esteem is related to self-compassion in women with female-pattern hair loss.

Research Method

In this research, the descriptive-correlation study method was used.

Statistical population and sample

The statistical population of this research included all women suffering from female pattern hair loss who visited the Ronic Skin and Hair Clinic in Qazvin city in the spring of 1402, and the statistical sample of the current study included 300 women suffering from female pattern hair loss. were selected as available sampling.

Tools

Self-esteem scale

The Rosenberg Self-Esteem Scale (RSES) (Rosenberg, 1965) was created in 1965. The scoring of this scale is based on a 4-point Likert scale, in which a score of 4 is given to the answer to completely agree and 1 is given to completely disagree. In the research (Greenberger et al, 2003), the reliability of this scale using the internal consistency method was 0.84 and the retest method was 0.84 after two weeks, and 0.67 after 5 months. In their research, it has also been stated that this scale has a significant negative relationship with the subscale of neuroticism and with the subscales of extroversion, empiricism, and duty-based from the Neo personality questionnaire, and not with the subscale of agreement, these results indicate validity. Convergent and divergent averages and suitable for this scale. In the research (Rajabi and Bahloul, 2016) for the reliability of this scale, the internal consistency coefficients of the items in the entire student sample were 0.84, male students 0.87, and female students 0.80. The correlation coefficients between each of the items of the scale with the total score of the items varied from 0.56 to 0.72 and all were significant at the $P < 0.001$ level. In their research, the analysis between the above scale and the death obsession scale showed -0.34 in the whole sample, -0.44 in male students, and -0.27 in female students, which indicates the divergent validity of Rosenberg's self-esteem scale.

Body Image Concern questionnaire

The Body Image Concern Questionnaire (BICI) was designed by (Littleton et al, 2005). The scoring of this questionnaire is based on a 5-point Likert scale so that the answer of never is given a score of 1 and always a score of 5. In this research, the reliability of this questionnaire was checked by the internal consistency method and Cronbach's alpha coefficient was 0.93. In their research, the convergent validity of this tool was checked by calculating its correlation coefficient with the Padova Obsessive Compulsive Questionnaire and the Eating Disorders Questionnaire, and its correlation coefficients with these tools were 0.62 and 0.40, respectively, in Mohammadi and Sajjadinejad's research. (2009) Cronbach's alpha coefficient of 0.84 and the result of factor analysis using the oblique rotation method and based on the Scree test, in total, all the questions explained 39.88% of the total variance of the test.

Compassion scale

The Self-Compassion Scale Short Form (SCS-SF) has 12 questions that were designed by (Raes et al, 2011). The scoring of this scale is using a 5-point Likert scale in such a way that a score of 1 is assigned to the answer "never" and "always" is assigned a score of 5. The reliability of this scale was reported to be 0.86 alpha coefficient and 0.97 correlation coefficient with the long form. In their research, after analyzing the confirmatory factor, they stated that the version of 12 questions is a suitable alternative to the long version of self-compassion. In Shahbazi et al.'s research (2014), the reliability of this scale using Cronbach's alpha was reported as 0.91 for the whole. In their research,

the concurrent validity coefficients of this scale with the general health questionnaire were significant from -0.28 to -0.48.

Data analysis methods

SPSS 26 software and descriptive and inferential tests were used to analyze the data. In descriptive statistics, frequency distribution, mean and standard deviation were used, and to test the research hypotheses, statistical tests of multiple regression analysis using stepwise method and Pearson's correlation coefficient were used.

Findings

Table No. 1. Descriptive statistics

The standard deviation	Average	the most	the least	Number	Variable
2/956	24/93	33	17	300	Self-esteem
8/413	56/78	79	37	300	Body image concerns
7/203	35/16	53	19	300	Self-compassion

According to table number 1, the average level of self-esteem in women with hair loss is 24.93, self-compassion is 35.16, and concern about body image is 56.78.

Table No. 2. Summary of self-esteem regression analysis

R ² Adjusted	R ²	R	Predictor variable	Model	Criterion variable
0/131	0/134	0/366	Body image concerns	1	Self-esteem
0/197	0/202	0/450	Body image concerns and self-compassion	2	

Table No. 2 shows that both predictive variables have been able to enter the equation and have a significant contribution in predicting self-esteem according to the amount of explained variances.

Table No. 3. Results of variance analysis of self-esteem

P	F	MS	df	SS	Sources Change	Model
0/000	46/117	350/043	1	350/043	regression	1
		7/590	298	2261/906	Remaining	
		-	299	2611/949	Total	
0/000	37/690	264/365	2	528/731	regression	2
		7/014	297	2083/219	Remaining	
		-	299	2611/949	Total	

Table No. 3 shows the results of the variance analysis of self-esteem according to the variables of body image concern and self-compassion. The significance levels of both models are less than 0.05 and therefore the explained variance ratios in each model are statistically significant.

Table No. 4. Regression coefficient of predictor variable in the model

P	t	Beta	SE	B	Predictive variables	Model
0/000	29/562	-	1/087	32/233	The constant of Eq	1
0/000	-6/791	-0/366	0/019	-0/129	Body image concerns	
0/000	20/341	-	1/366	27/790	The constant of Eq	2
0/000	-6/397	-0/334	0/018	-0/117	Body image concerns	
0/000	5/047	0/264	0/021	0/108	Self-compassion	

Table number 5. Pearson's correlation coefficient

Self-compassion	Body image concerns	statistics	Variable
0/304	-0/366	The correlation coefficient	Self-esteem
0/000	0/000	Significance level	
300	300	Number of subjects	

According to table number 5, the observed correlation coefficient for self-esteem and concern about body image is equal to -0.366 and the significance level is 0.000, and thus, with 95% confidence, we can say that there is a statistically significant relationship between self-esteem and concern about body image. There is. This relationship is inverse. On the other hand, the observed correlation coefficient for self-esteem and self-compassion is equal to 0.304 and the significance level is 0.000, and as a result, there is a statistically significant relationship between self-esteem and self-compassion, and this relationship is positive.

Discussion

According to what was mentioned, the following research was aimed at investigating the relationship between self-esteem, concern about body image and self-compassion in women suffering from hair loss with a female pattern. The female pattern was determined in the spring of 1402 when they referred to the Runic Skin and Hair Clinic in Qazvin city, and the statistical sample of the research included 300 women with female pattern hair loss who were selected through available sampling. Ronberg self-esteem scale (RSES), body image concern questionnaire (BICI) and self-compassion scale short form (SCS-SF) were used to collect data. After analyzing the scores, the results showed that self-esteem has a negative and statistically significant relationship with concern about body image in women with hair loss with a female pattern, and self-esteem with self-compassion in women with hair loss with a female pattern has a positive and statistically significant relationship. has it.

As the results showed, self-esteem and concern about body image in women with hair loss have a negative and statistically significant relationship with the female pattern. The result of this hypothesis is in line with the results of the following researches: (Liu et al., 2018) The results of the research mentioned that hair transplantation in men significantly increased the level of self-esteem and satisfaction with the appearance of patients. (Poot, 2004) stated in his research that patients with alopecia have lower self-esteem, quality of life and weaker body image. Self-esteem is closely related to appearance satisfaction (von Soest et al., 2009; Cash et al., 2002). The level of self-esteem is closely related to satisfaction with appearance, and many patients expect to increase their self-esteem through cosmetic surgery (Alfonso et al., 2005). Recently, studies have shown that cosmetic surgery not only improves physical appearance, but also increases self-esteem and builds self-confidence (von Soest et al., 2009; Cash et al., 2002).

To explain this finding from the research, it can be said that the body and dissatisfaction with it has become one of the most important concerns and preoccupations in the lives of girls and women (Ronoq Shashkalani et al., 2017). Hair is considered an important component of aesthetics for

women, hair loss with a female pattern also causes thinning and thinning of the hair, which reduces the beauty of the appearance of the hair to a part that can Dissatisfaction with the body image in women suffering from hair loss, and as a result, it causes them to withdraw from the society, and the continuation of this reduces their self-esteem, for another explanation of this research finding, this can be done. He stated that the intense worry of negative evaluation by others causes a decrease in satisfaction with the body image, and a person is faced with the question of whether my value is affected by my appearance and whether this appearance is acceptable. And this concern about body image continues to increase and causes a decrease in self-esteem in women with female pattern hair loss, and therefore self-esteem has a negative relationship with body image concern in women with female pattern hair loss.

On the other hand, the results showed that self-esteem and self-compassion in women with hair loss have a positive and statistically significant relationship with the female pattern, the result of this hypothesis is in line with the results of the following researches: the result of the research (Gandhi et al. 2023) showed that patients with alopecia areata experience psychological and psychosocial symptoms including depression, anxiety, anger, social withdrawal, embarrassment and low self-esteem. The research (Gilbert and Irons, 2005) on the relationship between self-esteem and compassion shows that positive self-esteem with positive value towards oneself helps to raise one's social position in interpersonal and social relationships, and the quality of compassion It increases kindness towards oneself. (Pisitsungkagarn et al, 2014) also showed that positive self-emotions are related to self-worth and better self-care.

In explaining this finding of the research, it can be stated as follows:

that people who have high self-compassion have a balanced and clear awareness of their experiences and hardly criticize themselves for their failures or not reaching their standards, and in fact ignore the painful aspects of their feelings or experiences taken and they do not ruminate (Neff and Vonk, 2009). In women with hair loss, when these women have self-compassion, they do not judge themselves too harshly for their hair loss, they do not use other people's feedback as a criterion for their own judgment, and problems arise. They accept their hair more easily and do not consider themselves worthless because of this, and they do not consider their value based on their appearance, on the contrary, if they have low compassion, they consider the components of their own and others' judgments about their appearance as proof of their value. They themselves know, so it can be concluded that self-esteem is related to self-compassion in women with female hair loss.

Offers

It is suggested to other researchers and researchers in their investigations and future studies in the society of women suffering from hair loss with a feminine pattern to investigate the relationship between external factors and variables and their effective role on self-esteem because it seems that these factors, along with other factors, can be a good predictor of self-esteem. Psychotherapists and counsellors are also suggested to have a special look at women suffering from female pattern hair loss, because so far not enough research has been done on this community and their psychological problems remain hidden, and since Women are psychologically more vulnerable than men, so more attention should be paid to the factors affecting their mental health.

References

1. Ayat-Elahi, Azin, Yazdan-Prest, Taraneh, Ahmad-Nasrollahi, Saman, Fatahi, Mehssa, Imam-Bakhsh, Mahdieh, Firouzeh, 1401, comparison of the effectiveness of topical solutions of vaproic acid 8.3% and minoxidil 5% in the treatment of hair loss. *Androgenetic hair. Skin and Beauty*, (13)1, 3-13.
2. Bayani, Azam, Pourhossein, Masoud and Bayani, Mohammad, Comparison of self-respect and irrational beliefs in women candidates and non-candidates for hair mesotherapy, 8th International Conference on Interdisciplinary Studies in Islamic and Human Sciences of Iran, August 1402, in press.
3. Rajabi, Gholamreza, Bahloul, Nasrin, 2016, reliability and validity assessment of Rosenberg self-esteem scale in first year students of Shahid Chamran University. *Journal of educational and psychological research*, third year, number 2, consecutive 8, autumn and winter.
4. Roanq Shashkalani, Mehrnaz, Panah Ali, Amir, Lotfi, Alireza, Qoimi, Mohammad Ali, Ahmedpour Pournaki, Farzin, 2017, comparison of body image and life satisfaction in cosmetic surgery applicants and those who underwent surgery in the maxillofacial and cosmetic surgery department of an educational hospital. *Treatment of Imam Reza of Tabriz. Razi Journal of Medical Sciences*, 25(8): 12-23.
5. Shahbazi, Massoud; Rajabi, Gholamreza; Megaki, Ibrahim; Jalodari, Arash, 2014, the confirmatory factor structure of the Persian version of the revised Shafqat - self-compassion rating scale in a group of prisoners, *psychological methods and models/6th year/19th issue*.
6. Mohammadi, Nurullah and Sajjadi-Nejad, Marzieh Al-Sadat, 2016, evaluation of psychometric indicators of body image concern questionnaire and test of correlation model of body mass index, dissatisfaction with body image and self-esteem in adolescent girls. *Journal of Psychological Studies, Faculty of Educational Sciences and Psychology, Al-Zahra University*, third volume, number 101, 1-85.
7. Adil, A., & Godwin, M. (2017). The effectiveness of treatments for androgenetic alopecia: a systematic review and meta-analysis. *Journal of the American Academy of Dermatology*, 77(1), 136-141.
8. Alessandrini, A, Bruni, F, Piraccini, BM, and Starace, M. Common causes of hair loss - clinical manifestations, trichoscopy and therapy. *J Eur Acad Dermatol Venereol*. (2021) 35:629–09. doi: 10.1111/jdv.17079.
9. Alfonso M, Richter-Appelt H, Tosti A, Viera MS, Garcia M. The psychosocial impact of hair loss among men: a multinational European study. *Curr Med Res Opin*. 2005; 21(11): 1829-1836.
10. Avis, N., Crawford, S., Manuel, J. (2004). Psychosocial problems among younger women with breast cancer. *Journal of Psycho-oncology*, 13, 295-308.
11. Banfield, S. & McCabe, M. (2002). An evaluation of the construct of body image. *Adolescence*, 37(146), 373-393.
12. Bertoli, M.J.; Sadoughifar, R.; Schwartz, R.A.; Lotti, T.M.; Janniger, C.K. Female pattern hair loss: A comprehensive review. *Dermatology* 2020, 33, e14055.
13. Cash TF, Duel LA, Perkins LL. Women's psychosocial outcomes of breast augmentation with silicone gel-filled implants: a 2-year prospective study. *Plast Reconstr Surg*. 2002; 109(6): 2112-2121.
14. Chaikittisilpa, S.; Rattanasirisin, N.; Panchaprateep, R.; Orprayoon, N.; Phutrakul, P.; Suwan, A.; Jaisamrarn, U. Prevalence of female pattern hair loss in postmenopausal women: A cross-sectional study. *Menopause* 2022, 29, 415–420.
15. Gandhi, K., Shy, M. E., Ray, M., Fridman, M., Vaghela, S., & Mostaghimi, A. (2023). The association of alopecia areata-related emotional symptoms with work productivity and daily activity among patients with alopecia areata. *Dermatology and Therapy*, 13(1), 285-298.
16. Gilbert P, Irons C. (2005). Focused therapies and com-passionate mind training for shame and self-attacking. *Compassion: Conceptualisations, research and use in psychotherapy*, 263-325.
17. Greenberger, E., Chen, C., Dmitrieva, J., & Farruggia, S.P. (2003). Item- wording and dimensionality of the Rosenberg self-esteem scale: Do they matter? *Personality and Individual Differences*, 35 (6), 12411254.
18. Greenberger, E., Chen, C., Dmitrieva, J., & Farruggia, S.P. (2003). Item- wording and dimensionality of the Rosenberg self-esteem scale: Do they matter? *Personality and Individual Differences*, 35 (6), 12411254.
19. Kranz D, Nadarevic L, Erdfelder E. Bald and Bad? *Exp Psychol*. 2019; 66(5): 331-345.
20. Littleton, H.L., Axsom, D.S. & Pury, C.L. (2005). Development of the body image concern inventory, *Behavior Research and Therapy*, 43:229 -241.
21. Liu, Fang; Miao, Yong; Li, Xingdong; Qu, Qian; Liu, Yang; Li, Kaitao; Feng, Chuanbo; Hu, Zhiqi (2018). The relationship between self-esteem and hair transplantation satisfaction in male androgenetic alopecia patients. *Journal of Cosmetic Dermatology* Volume 18, Issue 5 p. 1441-1447.
22. Masood, H. (2023). Efficacy of Scalp Cooling in the Prevention of Chemotherapy Induced Alopecia Among Breast Cancer Patients. *Sciential-McMaster Undergraduate Science Journal*, (10).
23. Müller Ramos, P., Melo, D. F., Radwanski, H., de Almeida, R. F. C., & Miot, H. A. (2023). Female-pattern hair loss: therapeutic update. *Anais Brasileiros de Dermatologia*, 98(4), 506-519.

24. Neff, K. D., Vonk R.(2009). Self-compassion versus global self-esteem:two different ways of relating to oneself. *Journal of Personality*,77(1),23 – 50.
25. Pisitsungkagarn, K., Taephant, N. and Attasaranya, P. (2014). Body image satisfaction and selfesteem in Thai female adolescents: the moderating role of selfcompassion. *International journal of adolescent medicine and health*, 26(3), 333-338.
26. Poot, F. (2004). Psychological consequences of chronic hair diseases. *Revue medicale de Bruxelles*, 25(4), A286-8.
27. Raes, F., Pommier, E., Neff, K. D., & Van Gucht, D. (2011). Construction and factorial validation of a Short Form of the Selfcompassion Scale. *Clinical Psychology and Psychotherapy*, 18, 250 255.
28. Ramos, P.M. Miot H.A. (2015). Female pattern hair loss: a clinical and pathophysiological review *An Bras Dermatol*, 90. pp. 529-543.
29. Ravary A, Baldwin MW. (2018). Self-esteem vulnerabilities are associated with cued attentional biases toward rejection. *Personality and Individual Differences*, 126, 44-51.
30. Rosenberg, M. (1965). *Society and the adolescent self- image*. Princeton, NJ: Princeton University Press.
31. Shimizu, G.K.M. Wedy G.F. , Schaefer L.V. Ramos, , P.M Miot. H.A. (2018), Translation into Portuguese language (Brazil), transcultural adaptation and validation of the quality of life questionnaire in female pattern hair loss (WAA-QoL-BP) *An Bras Dermatol*, 93 pp. 701-706.
32. Stapleton, P., Crighton, G. J., Carter, B. and Pidgeon, A. (2017). Self-Esteem and Body Image in Females: The Mediating Role of Self-Compassion and Appearance Contingent Self-Worth. *The Humanistic Psychologist*, 45(3): 238-257.
33. Suwan, A.; Jaisamrarn, U. Prevalence of female pattern hair loss in postmenopausal women: A cross-sectional study. *Menopause* 2022, 29, 415–420.
34. Tsutsui G.M. Ramos, P.M. Miot, H.A. (2022), Prevalence of female pattern hair loss in a multiracial population *J Am Acad Dermatol*, 86 pp. 862-894.
35. von Soest T, Kvalem IL, Roald HE, Skolleborg K. The effects of cosmetic surgery on body image, self-esteem, and psychological problems. *J Plast Reconstr Aesthet Surg*. 2009; 62(10): 1238-1244.
36. Youssef, S.M.E.; Atallah, R.B.; Zaky, M.S.; Eldeek, B.S.; Elsaie, M.L. Urban-rural differences in the prevalence of female pattern hair loss among secondary school girls: A cross-sectional study. *J. Cosmet. Dermatol*. 2022, 21, 2229–2235.